

Dyonic black holes in effective string theory

S. Mignemi

Dipartimento di Scienze Fisiche, Università di Cagliari
via Ospedale 72, 09100 Cagliari, Italy

The spherical symmetric dyonic black hole solutions of the effective action of heterotic string are studied perturbatively up to second order in the inverse string tension. An expression for the temperature in terms of the mass and the electric and magnetic charge of the black hole is derived and it is shown that its behaviour is qualitatively different in the two special cases where the electric or the magnetic charge vanishes.

February 1993

At present, string theories are the most promising attempt to unify gravitation with the other fundamental interactions. It is therefore of great interest to study their phenomenological consequences, in particular in the context of gravitational physics.

For this reason, the modifications to black hole physics induced by string theory have been extensively investigated. Due to the difficulty of treating the full theory, a fruitful approach has been the study of its low energy limit by means of effective field theoretic actions, obtained as an expansion in the inverse string tension α' . These actions describe the dynamics of the low energy excitations of the string spectrum, which comprise in the simplest case the graviton, a dilaton, an axion and gauge fields. In particular, some exact solutions have been obtained for the lowest order approximation to the action [1-3]. The main difference from the Einstein theory is the presence of a non-minimally coupling of the dilaton with the other fields, which changes drastically the physical properties of the black hole, and in particular the thermodynamics. For charged black holes, for example, the temperature becomes independent of the charge, at variance with the Reissner-Nordström solution of the Einstein-Maxwell theory.

The results obtained in [1-3] are however valid only if the order α' corrections in the gravitational sector of the effective action are neglected. These consist essentially in the presence of a higher derivative Gauss-Bonnet term coupled to the dilaton. In a recent paper [4], we have examined how the magnetic charged dilatonic black hole solutions of effective string theories obtained in [1] are modified when the order α' terms in the effective action are taken into account. Of particular interest were the thermodynamical properties of the solutions, which, contrary to the solutions of the leading order action, exhibit a dependence of the temperature on the magnetic charge of the black hole. In particular, these solutions can reach a vanishing temperature for a finite value of the mass, and therefore admit a stable remnant as final state of the Hawking evaporation.

In this paper we extend our investigation to the case of dyonic black holes. When higher order corrections to the action are neglected, the dyonic solution has been obtained by the magnetic one [1] by exploiting the $SL(2,R)$ dilaton-axion symmetry of the leading order terms in the perturbative expansion of the action [2]. Even if it has been conjectured that this symmetry might be an exact symmetry of the full theory [3], it is not however

a symmetry of the order α' action, since the Gauss-Bonnet term in that action is not invariant under the $SL(2, R)$ symmetry. It has been argued however that the symmetry could still be present in a highly non-trivial and non-linear way and therefore not order by order in perturbation theory [5].

For these reasons we perform here a perturbative calculation of the spherical symmetric solutions of the order α' action around the background constituted by the Schwarzschild solution. The most interesting result of our analysis is the difference in the behaviour of the temperature for an electrically charged black hole with respect to the magnetic one when higher order terms are taken into account. In fact, it appears that the corrections to the temperature have opposite sign in the two cases, so that for a purely electric black hole in this theory the temperature is a monotonic decreasing positive definite function of the mass for any value of the charge and therefore does not give rise to stable remnants.

The bosonic sector of the dimensionally reduced effective action for the heterotic string is given up to order α' by [6]:

$$S_{eff} = \int d^4x \sqrt{-g} \left[\mathcal{R} - \frac{1}{3} e^{-4\Phi} H_{abc} H_{abc} - 2(\nabla\Phi)^2 + \alpha e^{-2\Phi} (\mathcal{S} - F^2) \right] \quad (1)$$

where $\alpha \equiv \alpha'/8$, F is the Maxwell field strength, $\mathcal{S} \equiv \mathcal{R}_{abcd}^2 - 4\mathcal{R}_{ab}^2 + \mathcal{R}^2$ is the Gauss-Bonnet term and we have omitted the terms which do not contribute to the field equations up to order α^2 . The axion field H_{abc} is given explicitly in terms of the potential B_{ab} as:

$$H_{abc} = \partial_{[a} B_{bc]} + \alpha (A_{[a} \partial_b A_{c]} + \omega^{lm}_{[a} \partial_b \omega^{lm}_{c]} + \omega^{lm}_{[a} \omega^{mn}_b \omega^{nl}_{c]}),$$

where the Chern-Simons terms have been taken into account. Due to the presence of these terms a solution presenting both electric and magnetic charge will also have a non-trivial axion field [6].

It is well known that if one defines a scalar field a , dual to H , such that

$$H_{abc} = -\frac{1}{2} \epsilon_{abcd} e^{4\Phi} \partial_d a \quad (2)$$

the field equations can be derived from the action:

$$S_{eff} = \int d^4x \sqrt{-g} \left[\mathcal{R} - 2(\nabla\Phi)^2 - \frac{1}{2} e^{4\Phi} (\nabla a)^2 - \alpha e^{-2\Phi} (F^2 - \mathcal{S}) \right] - \alpha a (F\tilde{F} - \mathcal{R}\tilde{\mathcal{R}}) \quad (3)$$

where

$$F\tilde{F} = \frac{1}{2}\epsilon_{abcd}F_{ab}F_{cd} \quad \mathcal{R}\tilde{\mathcal{R}} = \frac{1}{2}\epsilon_{abcd}\mathcal{R}_{abef}\mathcal{R}_{cdef} \quad (4)$$

The field equations can then be written as:

$$\begin{aligned} \mathcal{R}_{mn} = & 2\nabla_m\Phi\nabla_n\Phi + \frac{1}{2}\nabla_m a\nabla_n a + 2\alpha e^{-2\Phi}(F_{mp}F_{np} - \frac{1}{4}g_{mn}F^2) \\ & + 4\alpha e^{-2\Phi}\left[4\mathcal{R}_{p(m}\nabla_{n)}\nabla_p\Phi - 2\mathcal{R}_{mn}\nabla_p\nabla_p\Phi - g_{mn}\mathcal{R}_{pq}\nabla_p\nabla_q\Phi \right. \\ & \left. - \mathcal{R}\nabla_m\nabla_n\Phi + \frac{1}{2}g_{mn}\mathcal{R}\nabla_p\nabla_p\Phi - 2\mathcal{R}_{qmn p}\nabla_p\nabla_q\Phi\right] \\ & - 8\alpha e^{-2\Phi}\left[4\mathcal{R}_{p(m}\nabla_{n)}\Phi\nabla_p\Phi - 2\mathcal{R}_{mn}\nabla_p\Phi\nabla_p\Phi - g_{mn}\mathcal{R}_{pq}\nabla_p\Phi\nabla_q\Phi \right. \\ & \left. - \mathcal{R}\nabla_m\Phi\nabla_n\Phi + \frac{1}{2}g_{mn}\mathcal{R}\nabla_p\Phi\nabla_p\Phi - 2\mathcal{R}_{qmn p}\nabla_p\Phi\nabla_q\Phi\right] \\ & + 4\alpha\left[\epsilon_{blmn}\mathcal{R}_{ajmn}\nabla_l\nabla_j a + \epsilon_{almn}\mathcal{R}_{bjmn}\nabla_l\nabla_j a\right] \end{aligned} \quad (5a)$$

$$\nabla^2\Phi = \frac{\alpha}{2}e^{-2\Phi}(\mathcal{S} - F^2) + \frac{1}{2}e^{4\Phi}(\nabla_p a)^2 \quad (5b)$$

$$\nabla_p(e^{4\Phi}\nabla_p a) = \alpha(F\tilde{F} - \mathcal{R}\tilde{\mathcal{R}}) \quad (5c)$$

$$\nabla_p(e^{-2\Phi}F_{pm}) = \frac{1}{2}\epsilon_{pmqr}F_{qr}\nabla_r a \quad (5d)$$

It is convenient to define a new scalar field b , such that

$$\nabla_p b = e^{4\Phi}\nabla_p a \quad (6)$$

The field equation can then easily be written in term of b . In particular:

$$\nabla^2\Phi = \frac{\alpha}{2}e^{-2\Phi}(\mathcal{S} - F^2) + \frac{1}{2}e^{-4\Phi}(\nabla b)^2 \quad (7a)$$

$$\nabla^2 b = \alpha(F\tilde{F} - \mathcal{R}\tilde{\mathcal{R}}) \quad (7b)$$

We want to find a perturbative expansion of the solution to (5) around the background constituted by the Schwarzschild metric of mass m with vanishing dilaton and axion. Our expansion will be in the parameter α or, more correctly, in $\frac{\alpha}{m^2}$. We adopt a spherically symmetric ansatz for the metric:

$$ds^2 = -\lambda^2 dt^2 + \lambda^{-2} dr^2 + R^2 d\Omega^2 \quad (8)$$

where $\lambda = \lambda(r)$, $R = R(r)$. This particular form of the metric is suggested from the exact solution found in [2] in the absence of the Gauss-Bonnet term (see also [4]).

The general spherical symmetric solution of the generalized Maxwell equation (5d) containing both electric and magnetic charge q_e and q_m respectively, is given in orthonormal coordinates by:

$$F_{ij} = \frac{q_m}{R^2} \quad F_{01} = \frac{q_e + q_m a}{R^2} e^{2\Phi} \quad (9)$$

We can now expand the fields in α as:

$$\begin{aligned} \lambda &= \lambda_0(1 + \alpha\psi_1 + \alpha^2\psi_2 + \dots) & R &= r + \alpha\rho_1 + \alpha^2\rho_2 + \dots \\ \Phi &= \alpha\phi_1 + \alpha^2\phi_2 + \dots & b &= \alpha\chi_1 + \alpha^2\chi_2 + \dots \end{aligned} \quad (10)$$

where $\lambda_0 = (1 - \frac{2m}{r})^{1/2}$ and ψ_i , ρ_i , ϕ_i and χ_i are functions of r . Moreover, from (9)

$$F_{ij} \sim \frac{q_m}{r^2} \left(1 - \frac{2\alpha\rho_1}{r} + \dots\right) \quad F_{01} \sim \frac{q_e}{r^2} + \alpha \left[\frac{q_e}{r^2} \left(2\phi_1 - \frac{2\rho_1}{r}\right) + \frac{q_m}{r^2} \chi_1 \right] + \dots \quad (11)$$

and hence:

$$F^2 \sim 2 \frac{q_m^2 - q_e^2}{r^4} + 2\alpha \left[-4q_m^2 \frac{\rho_1}{r^5} + 4q_e^2 \left(\frac{\rho_1}{r^5} - \frac{\phi_1}{r^4} \right) - 2q_e q_m \frac{\chi_1}{r^4} \right] + \dots \quad (12a)$$

$$F\tilde{F} \sim 4 \frac{q_e q_m}{r^4} + 4\alpha \left[q_m^2 \frac{\chi_1}{r^4} + 2q_e q_m \left(\frac{\phi_1}{r^4} - \frac{2\rho_1}{r^5} \right) \right] + \dots \quad (12b)$$

The field equations can now be expanded by inserting (10) and (11). At first order in α one has:

$$[r(r-2m)\phi_1']' = \frac{q_e^2 - q_m^2}{r^2} + \frac{24m^2}{r^4} \quad (13a)$$

$$[r(r-2m)\chi_1']' = 4 \frac{q_e q_m}{r^2} \quad (13b)$$

$$\rho_1'' = 0 \quad (13c)$$

$$[(r-2m)\psi_1]' = -\frac{m}{r^2}\rho_1 - \frac{q_e^2 + q_m^2}{2r^2} \quad (13d)$$

If one requires asymptotic flatness, a solution can be obtained such that $\psi_1 = 0$, namely:

$$\phi_1 = -\frac{1}{m} \left(\frac{2 + q_e^2 - q_m^2}{2r} + \frac{m}{r^2} + \frac{4m^2}{3r^3} \right), \quad \chi_1 = -\frac{2q_e q_m}{mr},$$

$$\rho_1 = -\frac{q_e^2 + q_m^2}{2m}, \quad \psi_1 = 0. \quad (14)$$

We pass now to evaluate the second order corrections, which satisfy the equations:

$$\begin{aligned} [r(r-2m)\phi_2']' &= 2\frac{r-2m}{r}\rho_1\phi_1' - \frac{48m^2}{r^4}\left(\phi_1 + \frac{\rho_1}{r}\right) + 2(q_m^2 + q_e^2)\frac{\phi_1}{r^2} \\ &\quad + 4(q_m^2 - q_e^2)\frac{\rho_1}{r^3} + 2q_e q_m \frac{\chi_1}{r^2} + \frac{r(r-2m)}{2}\chi_1'^2 \\ [r(r-2m)\chi_2']' &= 2\frac{r-2m}{r}\rho_1\chi_1' + 4q_m^2 \frac{\chi_1}{r^2} + 8q_m q_e \left(\frac{\phi_1}{r^2} - 2\frac{\rho_1}{r^3}\right) \\ \rho_2'' &= -r\phi_1'^2 + \frac{8m}{r^2}\phi_1'' - \frac{1}{4}r\chi_1'^2 \\ [(r-2m)\psi_2]' &= -\frac{r-2m}{2}\rho_2'' - \frac{r-m}{r}\rho_2' - \frac{m}{r^2}\rho_2 + \frac{2m}{r^3}\rho_1^2 - (q_e^2 - q_m^2)\frac{\phi_1}{r^2} \\ &\quad + 2(q_m^2 + q_e^2)\frac{\rho_1}{r^3} - 4m\left(\frac{r-2m}{r^2}\phi_1'' - 2\frac{r-3m}{r^3}\phi_1'\right) - q_e q_m \frac{\chi_1}{r^2} \end{aligned} \quad (15)$$

The asymptotically flat solutions of (15), with boundary conditions such that m is the physical mass of the black hole, are given by:

$$\begin{aligned} \phi_2 &= -\left[\frac{73-45q_m^2}{60m^3r} + \frac{73-15q_m^2+30q_e^2-15q_m^4-60q_m^2q_e^2+15q_e^4}{60m^2r^2}\right. \\ &\quad \left.+ \frac{73+45q_e^2}{45mr^3} + \frac{73+5q_m^2+50q_e^2}{30r^4} + \frac{112m}{75r^5} + \frac{8m^2}{9r^6}\right] \\ \chi_2 &= q_e q_m \left[\frac{3}{2m^3r} + \frac{3-2(q_m^2+q_e^2)}{2m^2r^2} + \frac{2}{3mr^3} + \frac{1}{3r^4}\right] \\ \rho_2 &= -\left[\frac{4-4(q_m^2-q_e^2)+(q_m^2+q_e^2)^2}{8m^2r} + \frac{2-q_m^2+q_e^2}{3mr^2} + \frac{7-3(q_m^2-q_e^2)}{3r^3} + \frac{16m}{5r^4} + \frac{24m^2}{5r^5}\right] \\ (r-2m)\psi_2 &= -\left[\frac{1-2(q_m^2-q_e^2)}{3mr^2} - \frac{11-5(q_m^2-q_e^2)^2}{3r^3} + \frac{(2-50(q_m^2-q_e^2))m}{15r^4} + \frac{272m^3}{15r^6}\right] \end{aligned} \quad (16)$$

and hence, up to order α^2 :

$$\begin{aligned} R^2 &\sim r^2 - \alpha \frac{q_m^2 + q_e^2}{m} r - \alpha^2 \frac{1 + q_e^2 - q_m^2}{m^2} + O\left(\frac{1}{r}\right) \\ \lambda^2 &\sim 1 - \frac{2m}{r} - \frac{1 + 2q_e^2 - 2q_m^2}{3mr^3} \alpha^2 + O\left(\frac{1}{r^4}\right) \\ \Phi &\sim -\left[\frac{2 + q_e^2 - q_m^2}{2m} \alpha + \frac{73 - 45q_m^2}{60m^3} \alpha^2\right] \left(\frac{1}{r}\right) + O\left(\frac{1}{r^2}\right) \\ a &\sim -q_e q_m \left[\left(\frac{2\alpha}{m} + \frac{3\alpha^2}{m^2}\right) \frac{1}{r} + \frac{5 + 6q_e^2 - 2q_m^2}{2m^2r^2} \alpha^2\right] + O\left(\frac{1}{r^2}\right) \end{aligned} \quad (17)$$

The black hole has therefore dilatonic and axionic charge given respectively at order α by $D = \frac{2+q_e^2-q_m^2}{2m}\alpha$ and $A = \frac{2q_e q_m}{m}\alpha$; these are not however independent parameters, but are functions of q_e , q_m and m , in accordance with the weak form of the no-hair conjecture [2]. The metric has a singularity at $r = r_- \sim \alpha \frac{q_e^2+q_m^2}{m}$ and a horizon at $r = r_+ \sim 2m \left(1 - \frac{1-2(q_e^2+q_m^2)^2}{12m^2}\alpha^2\right)$. We notice that when $r_- > r_+$, the singularity is naked. This regime is however out of the range of our approximations.

The temperature of the black hole can be readily obtained by requiring the regularity of the Euclidean section and is given by [4]:

$$T^{-1} \equiv \beta \sim 8\pi m \left[1 - \alpha^2 \left(\frac{\tilde{\psi}_2(2m)}{m} - 2\tilde{\psi}'_2(2m) \right) \right] = 8\pi m \left(1 - \alpha^2 \frac{73 - 45(q_m^2 - q_e^2)}{120m^4} \right) \quad (18)$$

This should be compared with the value $T = (8\pi m)^{-1}$ found in [2] when the order α' corrections to the action are neglected: the temperature is no longer independent of the electric and magnetic charge. However, while for a black hole carrying only magnetic charge the temperature has a maximum and then goes to zero for a finite value of m [4], the temperature of a purely electric black hole is at this order of approximation a monotonic decreasing function of the mass and therefore no massive remnant is to be expected in the last case.

Appendix

We compare here the perturbative result in the absence of higher order corrections with the exact solution found in [2]. If one neglects the Gauss-Bonnet term, the field equations are, at first order:

$$\begin{aligned} [r(r-2m)\phi'_1]' &= \frac{q_e^2 - q_m^2}{r^2} \\ [r(r-2m)\chi'_1]' &= 4\frac{q_e q_m}{r^2} \\ \rho''_1 &= 0 \\ [(r-2m)\psi_1]' &= -\frac{m}{r^2}\rho_1 - \frac{q_e^2 + q_m^2}{2r^2} \end{aligned} \quad (A.1)$$

whose solutions are:

$$\phi_1 = \frac{q_m^2 - q_e^2}{2mr}, \quad \chi_1 = -\frac{q_e q_m}{mr}, \quad \rho_1 = -\frac{q_e^2 + q_m^2}{2m}, \quad \psi_1 = 0 \quad (A.2)$$

At second order, the field equations become:

$$\begin{aligned}
[r(r-2m)\phi_2']' &= 2\frac{r-2m}{r}\rho_1\phi_1' + 2(q_m^2 + q_e^2)\frac{\phi_1}{r^2} + 4(q_m^2 - q_e^2)\frac{\rho_1}{r^3} \\
&\quad + 2q_e q_m \frac{\chi_1}{r^2} + \frac{1}{2}r(r-2m)\chi_1'^2 \\
[r(r-2m)\chi_2']' &= 2\frac{r-2m}{r}\rho_1\chi_1' + 4q_m^2 \frac{\chi_1}{r^2} + 8q_m q_e \left(\frac{\phi_1}{r^2} - 2\frac{\rho_1}{r^3} \right) \\
\rho_2'' &= -r\phi_1'^2 - \frac{1}{4}r\chi_1'^2 \\
[(r-2m)\psi_2]' &= -\frac{r-2m}{2}\rho_2'' - \frac{r-m}{r}\rho_2' - \frac{m}{r^2}\rho_2 + \frac{2m}{r^3}\rho_1^2 \\
&\quad - (q_e^2 - q_m^2)\frac{\phi_1}{r^2} + 2(q_m^2 + q_e^2)\frac{\rho_1}{r^3}
\end{aligned} \tag{A.3}$$

After a lengthy but straightforward calculation, one obtains:

$$\begin{aligned}
\phi_2 &= \frac{q_e^4 - 4q_e^2 q_m^2 - q_m^4}{4m^2 r^2}, \quad \chi_2 = -q_e q_m \frac{q_e^2 + q_m^2}{m^2 r^2} \\
\rho_2 &= -\frac{(q_e^2 + q_m^2)^2}{8m^2 r}, \quad \psi_2 = 0
\end{aligned} \tag{A.4}$$

Substituting the results in (10, 11) yields:

$$\begin{aligned}
\lambda^2 &= 1 - \frac{2m}{r} \quad R^2 = r^2 - \alpha \frac{(q_e^2 + q_m^2)}{m} r \quad F_{01} = \frac{q_e}{r^2} \\
\Phi &= -\left(\alpha \frac{q_e^2 - q_m^2}{2mr} + \alpha^2 \frac{q_e^4 - 4q_e^2 q_m^2 - q_m^4}{4m^2 r^2} \right) + \dots \quad a = -\alpha \frac{2q_e q_m}{mr^2} \left(1 + \alpha \frac{3q_e^2 - q_m^2}{2mr} \right) + \dots
\end{aligned}$$

This results coincide with the expansion in α of the exact solution found in (3):

$$\begin{aligned}
ds^2 &= -\left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r \left(r - \frac{\alpha(q_e^2 + q_m^2)}{2m} \right) d\Omega^2 \\
\Phi &= -\frac{1}{2} \ln \left(\frac{mr(mr - \alpha(q_e^2 - q_m^2))}{(mr)^2 - 2\alpha q_e^2 mr + \alpha^2 q_e^2 (q_e^2 + q_m^2)} \right) \\
a &= -q_e q_m \frac{2mr - \alpha(q_e^2 + q_m^2)}{(mr)^2 - 2\alpha q_e^2 mr + \alpha^2 q_e^2 (q_e^2 + q_m^2)} \\
F_{01} &= \frac{q_e}{r^2} \quad F_{23} = \frac{q_m}{r(r - \alpha(q_e + q_m)/2m)}
\end{aligned}$$

Acknowledgements

I wish to thank N.R. Stewart for many helpful discussions.

References

- [1] D. Garfinkle, G.T. Horowitz and A. Strominger, Phys. Rev. D43, 3140 (1991)
- [2] A. Shapere, S. Trivedi and F. Wilczek, Mod. Phys. Lett. A6, 2677 (1991)
- [3] A. Sen, "*Electric-magnetic duality in string theory*", preprint hep-th 9207053
- [4] S. Mignemi and N.R. Stewart, "*Charged black holes in effective string theory*", preprint hep-th 9212146
- [5] J.H. Schwarz, "*Dilaton-axion symmetry*", preprint hep-th 9209125
- [6] B.A. Campbell, M.J. Duncan, N. Kaloper and K.A. Olive, Nucl. Phys. B351, 778 (1991)